



Conserving wildlife and ecosystems focusing on protected areas and outstanding wild places





Sexual Selection, Temperature, and the Lion's Mane

Peyton M. West* and Craig Packer

The mane of the African lion (*Panthera leo*) is a highly variable trait that reflects male condition and ambient temperature. We examined the consequences of this variation in a long-term study of lions in the Serengeti National Park, Tanzania. Mane darkness indicates nutrition and testosterone and influences both female choice and male-male competition. Mane length signals fighting success and only appears to influence male-male assessment. Dark-maned males enjoy longer reproductive life-spans and higher offspring survival, but they suffer higher surface temperatures, abnormal sperm, and lower food intake during hot months of the year. Maned males are hotter than females, and males have lighter and/or shorter manes in hotter seasons, years, and habitats. This phenotypic plasticity suggests that the mane will respond to forecasted increases in ambient temperature.

Sexually selected indicator traits reflect male health and vigor (1), revealing how well individuals withstand environmental stress (2, 3). Environmental effects on trait morphology can be substantial (4, 5), outweighing both genetic effects (6) and reproductive advantages (7). Changing environmental conditions can alter trait costs, leading to the evolutionary loss of sexual ornaments (8) or possibly even to species extinction (9). The global environment is undergoing rapid

(10), and these changes have already altered some sexually selected behavior (11, 12). Sexually selected morphological traits may also be vulnerable, depending on the relative importance of ecological effects, reproductive benefits, and a trait's phenotypic and/or genetic plasticity, as well as the magnitude of environmental change.

change due to anthropogenic disturbance

Here we examine the evolutionary and ecological factors influencing the mane of the African lion. Manes are sexually dimorphic, develop at puberty, and are highly variable; thus, the mane has long been considered a sexually selected trait (13). Mane size and darkness are reduced in populations and subspecies living in hot climates [see supporting

Sustainable trophy hunting of African lions


Karyl Whitman, Anthony M. Starfield, Henley S. Quadling & Craig Packer

Department of Ecology, Evolution and Behavior, University of Minnesota, 1987 Upper Buford Circle, Saint Paul, Minnesota 55108, USA

In most species, sport hunting of male trophy animals can only reduce overall population size when the rate of removal of males is so high that females can no longer be impregnated¹. However, where males provide extensive paternal care, the removal of even a few individuals could harm the population as a whole^{2,3}. In species such as lions, excessive trophy hunting could theoretically cause male replacements (and associated infanticide^{4,5}) to become sufficiently common to prevent cubs reaching adulthood. Here we simulate the population consequences of lion trophy hunting using a spatially explicit, individual-based, stochastic model parameterized with 40 years of demographic data from northern Tanzania. Although our simulations confirm that infanticide increases the risk of population extinction, trophy hunting could be sustained simply by hunting males above a minimum age threshold, and this strategy maximizes both the

Department of Ecology, Evolution and Behavior, University of Minnesota, 1987 Upper Buford Circle, St. Paul. MN 55108. USA.

^{*}To whom correspondence should be addressed. E-mail: west0302@umn.edu





- **About FZS**
- The Biodiversity Crisis
- Protected Areas and Buffer Zones in Africa 3
- Perspective from the Ground 4
- **Moving Forward**

Serengeti Shall Not Die

FZS is one of Germany's oldest conservation organizations, active for more than 50 years in Serengeti and beyond






How FZS works

Biodiversity and wilderness

- Protected areas
- Partnerships
- Resource protection
- Community engagement








1	About FZS
2	The Biodiversity Crisis
3	Protected Areas and Buffer Zones in Africa
4	Perspective from the Ground
5	Moving Forward

"The biodiversity crisis (i.e. the rapid loss of species and the rapid degradation of ecosystems) is probably a greater threat than global climate change to the stability and prosperous future of humankind on Earth."

Professor Carsten Rahbek, Director for the Center for Macroecology, Evolution and Climate, University of Copenhagen

Planetary Boundaries

A safe operating space for humanity



Red

Beyond zone of uncertainty (high risk)

Yellow

In zone of uncertainty (increasing risk)

Green

Below boundary (safe)


"The main factor currently driving biodiversity loss is habitat destruction."


- Harvard University Center for Health and Global Environment





Protected areas in Africa





Threats to African Protected Areas










Buffer zone:

- An area peripheral to a national park or equivalent reserve, where restrictions are placed upon resource use or special development measures are undertaken to enhance the conservation values of the area.
- A gradient of protection around the core site

Source: UNEP-WCMC 2014, Biodiversity A-Z website: www.biodiversitya-z.org, UNEP-WCMC, Cambridge, UK.





- **About FZS**
- The Biodiversity Crisis
- Protected Areas and Buffer Zones in Africa 3
- Perspective from the Ground 4
- **Moving Forward**





Serengeti National Park, Tanzania





















Tanzania Prepares to
Hand Wildlife Reserves
Over to Farmers and
Livestock

SUSTAINABLE CITIES | RENEWABLE ENERGIES | WATER & ENVIRONMENT |

Homepage » TANZANIA: Five game reserves officially transformed into national parks

TANZANIA: Five game reserves officially transformed into national parks

By Jean Marie Takouleu - Published on February 28 2019 / Modified on February 28 2019





The Tanzanian government's plan to transform five game reserves into national parks was recently unanimously approved by parliament. Biharamuro, Burigi, Kimisi, Ibanda and Rumanyika thus become national parks.




North Luangwa National Park, Zambia











Gonarezhou National Park, Zimbabwe





14.03.20 © FZS

Gonarezhou elephant movement







- 1 About FZS
- 2 The Biodiversity Crisis
- 3 Protected Areas and Buffer Zones in Africa
- 4 Perspective from the Ground
- 5 Moving Forward



Protected Areas Work


theguardian

Biodiversity greater inside Earth's protected areas, study finds

Scientists show for the first time that there are 15% more individual plants and animals and 11% more species inside terrestrial conservation zones



A butterfly on a leaf in Yasuni national park, Ecuador, one of the world's biodiversity hotspots. Photograph:
 Dolores Ochoa/AP




Protected Area Spending Need in Developing Countries



Sources: James et al. BioScience, Volume 51, Issue 1, 1 January 2001, Pages 43–52; internal analysis



14.03.201 © FZS -







14.03.2019 © Amy Dickman



"From year to year, the threat of extermination hangs over these wild animals, and it is very unlikely that our children will ever see them as we can see them now—the vast procession of zebras, gazelles and wildebeest moving across the open plains..."

- Alan Morehead, 1960, Serengeti Shall Not Die

